

# UNCLASSIFIED

AD NUMBER
AD432596
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative Use; Feb 1964. Other requests shall be referred to U.S. Army Biological Laboratories, Fort Detrick, MD 21701.
AUTHORITY
ABL D/A ltr, 27 Sep 1971

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED  
AD **432596**

---

DEFENSE DOCUMENTATION CENTER  
FOR  
SCIENTIFIC AND TECHNICAL INFORMATION  
CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

432596

CATALOGED BY DDIC

AS AD No. \_\_\_\_\_

432596

TECHNICAL MANUSCRIPT 127

PLAQUE FORMATION  
BY PSITTACOSIS VIRUS

FEBRUARY 1964

MAR 20 1964

UNITED STATES ARMY  
BIOLOGICAL LABORATORIES  
FORT DETRICK

NO OTS

U.S. ARMY BIOLOGICAL LABORATORIES  
Fort Detrick, Frederick, Maryland

TECHNICAL MANUSCRIPT 127

PLAQUE FORMATION BY PSITTACOSIS VIRUS

Edmund H. Kozikowski

Nicholas Mahon

Aerobiology Division  
DIRECTOR OF BIOLOGICAL RESEARCH

Project IC022301A070

February 1964

A portion of the work reported here was performed under Project 4B11-02-068, "Aerobiological Research," Task -01, "Stability and Virulence of BW Aerosols." Expenditure orders were 2201301 and 2024. This material was originally submitted as manuscript 5312.

The information in this document has not been cleared for release to the public.

DDC AVAILABILITY NOTICE

Qualified requestors may obtain copies of this document from DDC.

Foreign announcement and dissemination of this document by DDC is limited.

PLAQUE FORMATION BY PSITTACOSIS VIRUS

ABSTRACT

The production of plaques by psittacosis virus on monolayers of McCoy cells is described in this preliminary report. Plaques appeared in five to seven days and ranged in diameter from 0.2 to 0.5 millimeter. Formation of plaques was inhibited by psittacosis antiserum, which confirmed their virus specificity. The relationship between plaque numbers and virus concentration was linear.

Although the production of plaques on cell monolayers has been reported for a considerable number of viral agents,<sup>1</sup> it has been demonstrated for only one member of the psittacosis group, meningeopneumonitis virus.<sup>2,3</sup> This preliminary communication describes plaque formation by another member, psittacosis (Borg) virus.

Cultures of an established cell line, McCoy, originally derived from human synovial tissue<sup>4</sup> and shown to be 99 per cent susceptible to infection with psittacosis virus,<sup>5</sup> were suspended in nutrient medium composed of mixture 199 with 0.5 per cent lactalbumin hydrolyzate (LAH), 10 per cent heat-inactivated calf serum, 50 micrograms of streptomycin, and 75 micrograms of kanamycin. Five milliliters of suspension containing  $1.5 \times 10^5$  cells was dispensed into two-ounce Brockway bottles (Sani-Glas, Brockway Glass Co.), capped, and incubated at 35°C. Continuous cell monolayers were formed, usually within 48 hours. These were washed once with ten milliliters of phosphate-buffered saline (PBS) and once with five milliliters of maintenance medium (mixture 199 with 5 per cent calf serum). A virus suspension that had been passed five times previously on McCoy cell cultures was employed in 0.5-milliliter volumes to infect cell monolayers. After an adsorption time of four hours at 35°C, residual inoculum was removed. Four milliliters of primary overlay medium, heated to 45°C, was added and cell monolayers were incubated four days at 35°C. The overlay medium contained one part calf serum, ten parts 2.2 per cent Noble agar (Difco), and ten parts nutrient medium. The latter consisted of one per cent LAH, 0.2 per cent yeast extract, and 0.04 per cent streptomycin in twofold concentrated Earle's salt solution. One milliliter of second overlay medium, differing only from primary overlay medium by the addition of 0.015 per cent neutral red, was added onto cell monolayers that were incubated an additional one to three days at 35°C.

On the fifth day, but more commonly on the sixth day, after cell monolayers were exposed to virus suspension, minute macroscopic plaques appeared as clear discrete areas ranging in diameter from 0.2 to 0.5 millimeter (Figure 1). Plaques examined microscopically consisted of unstained cells in a degenerate condition surrounded by a well-defined periphery of healthy stained cells. Although the size of plaques remained constant on subsequent incubation, their number increased gradually and reached a maximum on the seventh day. Enumeration was facilitated by employing a dissecting microscope at magnification 15X. By the tenth day, plaques tended to fade because of cell degeneration. Plaques produced by psittacosis virus were similar in size and appearance to those described for meningopneumonitis virus on strain L cell monolayers.<sup>3</sup> The maximum diameters of plaques produced by meningopneumonitis virus was attained on the eleventh day,<sup>3</sup> which suggests a difference between the rates of plaque formation by the two viruses. This may be related, however, to the use of different cell lines.

Plaque formation by different concentrations of virus was linear over a dilution range of 1.5 log units (Figure 2), suggesting that each plaque was initiated by a single virus particle. Significant reductions in plaque formation by varied dilutions of rooster psittacosis antiserum mixed with a standard suspension of virus confirmed the virus specificity of plaques.

In sensitivity, the plaque assay compared favorably with another technique, that of yolk-sac inoculation of seven-day chick embryonated eggs, based on dilution end point. Mean values of replicate determinations made by both techniques with a standard virus suspension were  $2.7 \times 10^5$  plaque-forming units (PFU) and  $2.8 \times 10^5$  egg LD<sub>50</sub> per milliliter, respectively. The feasibility of employing a plaque assay for psittacosis virus is indicated by these preliminary findings.






Figure 1. Plaques Formed by Psittacosis Virus on McCoy Cell Monolayers Six Days after Infection; Magnification 1X.




Figure 2. Linear Function Between Number of Plaques Formed on McCoy Cell Monolayers and Concentration of Psittacosis Virus. Each point is the mean of three determinations.

LITERATURE CITED

1. Cooper, P.D. "The plaque assay of animal viruses," In: Advances in virus research," New York, Academic Press, 1961. pp.319-378.
2. Tamura, A., and Higashi, N. "Plaque formation in monolayers of strain L cells infected with the meningopneumonitis virus," Ann. Rpt. Inst. Virus Res. Kyoto Univ. 2:57-64, 1959.
3. Higashi, N., and Tamura, A. "A plaque assay for meningopneumonitis virus in monolayers of strain L cells," Virology 12:578-588, 1960.
4. Fernandes, M.V. "Irradiation of cells in tissue culture. VII. Studies on the susceptibility to bluetongue virus of radiation-induced giant cells in vitro," Z. Zellforsch. Mikroskop. Anat. 50:433-443, 1959.
5. Hahon, N., and Nakamura, R.M. "Quantitative assay of psittacosis virus by the fluorescent cell-counting technique," Aerobiology Division, U.S. Army Biological Laboratories, Frederick, Maryland. December 1963. (Technical Manuscript 115).
6. Dulbecco, R. "Production of plaques in monolayer tissue cultures by single particles of an animal virus," Proc. Natl. Acad. Sci. U.S. 38:747-752, 1952.